



# CODING AND BILLING GUIDE FOR THE USE OF SOLIRIS® (eculizumab)

In Adult Patients With Anti-Aquaporin-4  
(AQP4) Antibody-Positive Neuromyelitis  
Optica Spectrum Disorder (NMOSD)

## INDICATION

SOLIRIS is indicated for the treatment of neuromyelitis optica spectrum disorder (NMOSD) in adult patients who are anti-aquaporin-4 (AQP4) antibody positive.

## SELECT IMPORTANT SAFETY INFORMATION

### WARNING: SERIOUS MENINGOCOCCAL INFECTIONS

SOLIRIS, a complement inhibitor, increases the risk of serious infections caused by *Neisseria meningitidis* [see *Warnings and Precautions* (5.1)]. Life-threatening and fatal meningococcal infections have occurred in patients treated with complement inhibitors. These infections may become rapidly life-threatening or fatal if not recognized and treated early.

- Complete or update vaccination for meningococcal bacteria (for serogroups A, C, W, Y, and B) at least 2 weeks prior to the first dose of SOLIRIS, unless the risks of delaying SOLIRIS therapy outweigh the risk of developing a serious infection. Comply with the most current Advisory Committee on Immunization Practices (ACIP) recommendations for vaccinations against meningococcal bacteria in patients receiving a complement inhibitor. See *Warnings and Precautions* (5.1) for additional guidance on the management of the risk of serious infections caused by meningococcal bacteria.
- Patients receiving SOLIRIS are at increased risk for invasive disease caused by *Neisseria meningitidis*, even if they develop antibodies following vaccination. Monitor patients for early signs and symptoms of serious meningococcal infections and evaluate immediately if infection is suspected.

Because of the risk of serious meningococcal infections, SOLIRIS is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called ULTOMIRIS and SOLIRIS REMS [see *Warnings and Precautions* (5.2)].

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

**SOLIRIS®**  
(eculizumab)  
Injection for Intravenous Use  
300 mg/30 mL vial

## Purpose of This Guide

Alexion Pharmaceuticals, Inc. has developed the SOLIRIS® (eculizumab) Coding and Billing Guide to provide objective and publicly available coding and billing information.

This document is provided for informational purposes only and is not legal advice or official guidance from payers. It is not intended to increase or maximize reimbursement by any payer. Alexion does not warrant, promise, guarantee, or make any statement that the use of this information will result in coverage or payment for SOLIRIS, or that any payment received will cover providers' costs. Alexion is not responsible for any action providers take in billing for, or appealing, SOLIRIS claims.

Hospitals and physicians are responsible for compliance with Medicare and other payer rules and requirements and for the information submitted with all claims and appeals. Before any claims or appeals are submitted, hospitals and physicians should review official payer instructions and requirements, should confirm the accuracy of their coding or billing practices with these payers, and should use independent judgment when selecting codes that most appropriately describe the services or supplies provided to a patient.

Please visit [www.SOLIRIS.net](http://www.SOLIRIS.net) for additional information or call 1-888-765-4747 to speak with the Alexion OneSource™ Team.

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

# Coding for SOLIRIS® (eculizumab) in anti-AQP4 antibody-positive NMOSD

## Diagnosis Coding

The following *International Classification of Diseases, 10th Revision, Clinical Modification* (ICD-10-CM) codes may be appropriate to describe adult patients diagnosed with anti-AQP4 antibody-positive NMOSD:

ICD-10-CM Diagnosis Code <sup>1</sup>	Code Descriptor
G36.0	Neuromyelitis optica [Devic]

## Drug Coding

The following drug-specific Healthcare Common Procedure Coding System (HCPCS) billing code can be reported on SOLIRIS medical claim forms to payers:

New HCPCS Code <sup>2</sup>	New Code Description	Effective for Dates of Service
<b>J1299</b>	<b>Injection, eculizumab, 2 mg</b>	<b>On and after April 1, 2025</b>
J1300	Injection, eculizumab, 10 mg	From January 1, 2008, through March 31, 2025

Amount of SOLIRIS Administered	300 mg	600 mg	900 mg	1200 mg	Effective Dates
<b>J1299 Billing Units</b>	150	300	450	600	<b>Starting April 1, 2025</b>
J1300 Billing Units	30	60	90	120	Through March 31, 2025

**1 billing unit of J1299 = 2 mg of SOLIRIS**  
**Billing with incorrect units may result in claim processing delays**

The following HCPCS modifiers may be required for SOLIRIS, as applicable:

Modifier <sup>2</sup>	Description	Commercial Requirement	Medicare Requirement
JZ	Zero drug amount discarded/not administered to any patient	Varies by payer	Y
RE	Furnished in full compliance with FDA-mandated risk evaluation and mitigation strategy (REMS)	Y	Y
TB	Drug or biological acquired with 340B drug pricing program discount, reported for informational purposes	N	Y

Some payers, including Medicaid, require drugs like SOLIRIS to be billed on medical claims with the product's National Drug Code (NDC) in addition to the HCPCS code. Payers typically require healthcare professionals to use the Health Insurance Portability and Accountability Act (HIPAA)-compliant, 11-digit NDC format<sup>4</sup>:

11-Digit NDC <sup>3,4</sup>	Code Descriptor	Strength
25682-0001-01	SOLIRIS single-use vial	300 mg/30 mL

Please note that payers have different guidance for placement of the NDC on medical claims. Typically, the 11-digit NDC is reported without any dashes or other punctuation.<sup>4</sup>

Some payers may also require a unit of measure (UoM) qualifier and quantity. For SOLIRIS, the UoM qualifier is mL (milliliter).

Check payer requirements for reporting the NDC and UoM on claims.

**Please note:** Changes to the HCPCS code and billing units do not affect NDC reporting.

Please see Important Safety Information on pages **1** and **11-12** and accompanying full **Prescribing Information** for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

## Drug Administration Services

The following are possible International Classification of Diseases, 10th Revision, Procedure Coding System (ICD-10-PCS) procedure codes to report the administration of SOLIRIS® (eculizumab) in inpatient settings:

ICD-10-PCS <sup>5</sup>	Code Descriptor
XW033C6	Introduction of eculizumab into peripheral vein, percutaneous approach, new technology group 6
XW043C6	Introduction of eculizumab into central vein, percutaneous approach, new technology group 6

The following Current Procedural Terminology (CPT®) codes may be appropriate to report administration of SOLIRIS in physician offices and hospital outpatient facilities. Individual payer policies should be reviewed for reporting requirements.

CPT <sup>6</sup>	Code Descriptor
96365	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); initial, up to one hour
+ 96366	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); each additional hour (list separately in addition to primary procedure)
96413 <sup>a</sup>	Chemotherapy administration, intravenous infusion technique; up to one hour, single or initial substance/drug
+ 96415 <sup>a</sup>	Chemotherapy administration, intravenous infusion technique; each additional hour (list separately in addition to primary procedure)

a. Billing highly complex administration codes (96413 and 96415) requires medical record documentation of the complexity involved beyond what is required for therapeutic infusions (96365 and 96366).<sup>7</sup>

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

# Coding for Meningococcal Vaccination

## Diagnosis Coding

For an encounter strictly for the vaccination, the diagnosis code for prophylactic vaccination is assigned along with the diagnosis code for NMOSD and any other conditions the patient may have.

ICD-10-CM Diagnosis Code <sup>1</sup>	Code Descriptor
Z23	Encounter for immunization

## Vaccine Coding

Coverage of meningococcal vaccines may vary by payer. Prescribers should consult respective payer billing guidelines.

CPT Code <sup>6</sup>	Code Descriptor
90619	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, tetanus toxoid carrier (MenACWY-TT), for intramuscular use
90620	Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2-dose schedule, for intramuscular use
90621	Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2- or 3-dose schedule, for intramuscular use
90623	Meningococcal pentavalent vaccine, conjugated Men A, C, W, Y-tetanus toxoid carrier, and Men B-FHbp, for intramuscular use
90624	Meningococcal pentavalent vaccine, Men B-4C recombinant proteins and outer membrane vesicle and conjugated Men A, C, W, Y-diphtheria toxoid carrier, for intramuscular use
90733	Meningococcal polysaccharide vaccine, serogroups A, C, Y, W-135, quadrivalent (MPSV4), for subcutaneous use
90734	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use
90749	Unlisted vaccine/toxoid

## Vaccine Administration Coding

The following CPT codes may be appropriate to report administration of meningococcal vaccines in outpatient settings:

CPT Code <sup>6</sup>	Code Descriptor
90471	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); 1 vaccine (single or combination vaccine/toxoid)
+ 90472	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure)

Please see Important Safety Information on pages 1 and 11-12 and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

# Claim Forms

## Sample CMS-1500 (or the electronic equivalent 837P): Physician Office

Example claim form for a SOLIRIS® (eculizumab) IV infusion:

To achieve a SOLIRIS maintenance dose of 1200 mg for a patient 18 years of age and older, the following vials were used:

- 4 single-use 300 mg/30 mL vials (NDC 25682-0001-01)

### Example claim for dates of service through March 31, 2025

21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E) ICD Ind.										22. RESUBMISSION CODE		ORIGINAL REF. NO.	
A. <b>G36.0</b> B. C. D.													
E. F. G. H.													
I. J. K. L.													
24. A. DATE(S) OF SERVICE From To B. PLACE OF SERVICE C. EMG D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) E. DIAGNOSIS POINTER F. \$ CHARGES G. DAYS OR UNITS H. EPSON Family Plan I. ID. QUAL. J. RENDERING PROVIDER ID. #													
1 N425682000101 ML120													
MM DD YY MM DD YY 11										J1300 JZ RE		A XXX XX 120 NPI	
2 MM DD YY MM DD YY 11										96365		A XXX XX 1 NPI	
3												NPI	

### Example claim for dates of service starting April 1, 2025

21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E) ICD Ind.										22. RESUBMISSION CODE		ORIGINAL REF. NO.	
A. <b>G36.0</b> B. C. D.													
E. F. G. H.													
I. J. K. L.													
24. A. DATE(S) OF SERVICE From To B. PLACE OF SERVICE C. EMG D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) E. DIAGNOSIS POINTER F. \$ CHARGES G. DAYS OR UNITS H. EPSON Family Plan I. ID. QUAL. J. RENDERING PROVIDER ID. #													
1 N425682000101 ML120													
MM DD YY MM DD YY 11										J1299 JZ RE		A XXX XX 600 NPI	
2 MM DD YY MM DD YY 11										96365		A XXX XX 1 NPI	
3												NPI	

Please see Important Safety Information on pages **1** and **11-12** and accompanying full **Prescribing Information** for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

# Claim Forms (for dates of service starting April 1, 2025)

## Sample CMS-1500 (or the electronic equivalent 837P): Physician Office

For an example of a completed CMS-1500 form, go to page [6](#).

**Item 21 Diagnosis:** Enter the appropriate diagnosis code; eg,  
– ICD-10-CM **G36.0** for neuromyelitis optica [Devic]

*Note: Other diagnosis codes may apply.*

**Item 23 Prior Authorization:** Enter the prior authorization number as obtained prior to services rendered.

☐ YES ☐ NO

22. RESUBMISSION CODE ORIGINAL REF. NO.

23. PRIOR AUTHORIZATION NUMBER

21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E)		ICD Ind.	
A. L	B. L	C. L	D. L
E. L	F. L	G. L	H. L
I. L	J. L	K. L	L. L
1			
2			
3			
4			

24. A. DATE(S) OF SERVICE	B. PLACE OF SERVICE	C. EMG	D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances)	E. DIAGNOSIS POINTER	F. \$ CHARGES	G. DAYS OR UNITS	H. PSUT Family Plan	I. ID. QUAL.	J. RENDERING PROVIDER ID. #
From To MM DD YY MM DD YY			CPT/HCPCS MODIFIER						
								NPI	
								NPI	
								NPI	
								NPI	

**Item 24A Date(s) of Service:** The “N4” qualifier is required before the NDC; do not include dashes.

Some payers may also require a UoM for each NDC; eg,

– N425682000101 ML30

*Note: Check payer requirements and format for reporting the NDC and UoM.*

**Item 24E Diagnosis Pointer:** Enter the letter (A-L) that corresponds to the diagnosis in Item 21.

**Item 24G Days or Units:** Enter the appropriate number of units of service; eg, SOLIRIS® (eculizumab) 1200 mg is reported with “600” units.

**Item 24D Procedures/Services/Supplies:** Enter the appropriate CPT/HCPCS codes and modifiers, eg,

- Drug: **J1299** Injection, eculizumab, 2 mg
- Applicable modifiers:
  - **JZ** Zero drug amount discarded/not administered to any patient
  - **RE** Furnished in full compliance with FDA-mandated REMS
- Administration: **96365** for IV infusion

*Note: Some payers may provide specific guidance.*

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

## Claim Forms

### Sample CMS-1450 (UB-04) (or the electronic equivalent 837I): Hospital Outpatient Clinic or Facility

Example claim form for a SOLIRIS® (eculizumab) IV infusion:

To achieve a SOLIRIS maintenance dose of 1200 mg for a patient 18 years of age and older, the following vials were used:

- 4 single-use 300 mg/30 mL vials (NDC 25682-0001-01)

#### Example claim for dates of service through March 31, 2025

42 REV. CD.	43 DESCRIPTION	44 HCPCS / RATE / HIPPS CODE	45 SERV. DATE	46 SERV. UNITS	47 TOTAL CHARGES
0636	N425682000101 ML120 SOLIRIS	J1300 JZ RE	MM DD YY	120	XXX XX
0510	IV infusion, initial, up to 1 hour	96365	MM DD YY	1	XXX XX

#### Example claim for dates of service starting April 1, 2025

42 REV. CD.	43 DESCRIPTION	44 HCPCS / RATE / HIPPS CODE	45 SERV. DATE	46 SERV. UNITS	47 TOTAL CHARGES
0636	N425682000101 ML120 SOLIRIS	J1299 JZ RE	MM DD YY	600	XXX XX
0510	IV infusion, initial, up to 1 hour	96365	MM DD YY	1	XXX XX

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.



## Claim Forms (for dates of service starting April 1, 2025)

### Sample CMS-1450 (UB-04) (or the electronic equivalent 837I): Hospital Outpatient Clinic or Facility

For an example of a completed CMS-1450 form, go to page [8](#).

#### Form Locator 42-43 Revenue Code and Description:

Enter the appropriate revenue code and description corresponding to the HCPCS code in field 44; eg,

- 0636 for drugs requiring detailed coding
- 0510 for clinic, general

*Note: Other revenue codes may apply.*

**Form Locator 46 Units of Service:** Enter the appropriate number of units of service; eg, SOLIRIS® (eculizumab) 1200 mg is reported with “600” units.

42 REV. CD.	43 DESCRIPTION	44 HCPCS / RATE / HIPPS CODE	45 SERV. DATE	46 SERV. UNITS	47 TOTAL CHARGES	48 NON-COVERED CHARGES
1						
2						
3						
4						
5						
6						
7						
8						

#### Form Locator 43 Revenue Description:

The “N4” qualifier is required before the NDC; do not include dashes.  
Some payers may require a UoM for each NDC; eg,

- N425682000101 ML30

*Note: Double check payer requirements and format for reporting the UoM.*

**Form Locator 44 HCPCS Code:** Enter the appropriate CPT/HCPCS codes and modifiers; eg,

- Drug: **J1299** Injection, eculizumab, 2 mg
- Applicable modifiers:
  - **JZ** Zero drug amount discarded/not administered to any patient
  - **RE** Furnished in full compliance with FDA-mandated REMS
- Administration: **96365** for IV infusion

*Note: Some payers may provide specific guidance.*

50 PAYER NAME		51 HEALTH PLAN ID	52 REL. INFO	53 ASG. BEN.	54 PRIOR PAYMENTS	55 EST. AMOUNT DUE	56 NPI	57 OTHER PRV ID	62 INSURANCE GROUP NO.
A									
B									
C									
58 IN.									62 INSURANCE GROUP NO.
A									
B									
C									
63 TR									ME
A									
B									
C									
66 DX									68
A									
B									
C									
69 ADMIT DX		70 PATIENT REASON DX		71 PPS CODE		72 ECI		73	
a.		b.		c.		d.		e.	
74 PRINCIPAL PROCEDURE CODE		75 OTHER PROCEDURE CODE		76 ATTENDING NPI		77 QUAL		78	


#### Form Locator 67 Principal Diagnosis Code and 67A-67Q Other Diagnosis Codes:

Enter the appropriate diagnosis code; eg,

- ICD-10-CM **G36.0** for neuromyelitis optica [Devic]

*Note: Other diagnosis codes may apply.*

Please see Important Safety Information on pages [1](#) and [11-12](#) and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.



## ALEXION ACCESS NAVIGATOR

Alexion Access Navigator is a dedicated resource website for US Healthcare Professionals and their offices that contains downloadable access and reimbursement materials for SOLIRIS® (eculizumab).

Online: <https://alexionaccessnavigator.com>

## OneSource™ Offers Patient Support

Contact OneSource:

**Phone:**  
1-888-765-4747

**Online:**  
<https://alexiononesource.com>

## References

1. Centers for Medicare & Medicaid Services. 2025 ICD-10-CM. Accessed February 21, 2025. <https://www.cms.gov/medicare/coding-billing/icd-10-codes#>
2. Centers for Medicare & Medicaid Services. April 2025 alpha-numeric HCPCS file. Accessed March 4, 2025. <https://www.cms.gov/medicare/coding-billing/healthcare-common-procedure-system/quarterly-update>
3. SOLIRIS. Prescribing Information. Alexion Pharmaceuticals, Inc.
4. Food and Drug Administration. Future format of the National Drug Code; public hearing; request for comments. *Fed Regist.* 2018;83(152):38666-38668. Accessed February 21, 2025. <https://www.federalregister.gov/documents/2018/08/07/2018-16807/future-format-of-the-national-drug-code-public-hearing-request-for-comments>
5. Centers for Medicare & Medicaid Services. 2025 ICD-10-PCS conversion table. Accessed February 21, 2025. <https://www.cms.gov/medicare/coding-billing/icd-10-codes>
6. American Medical Association. *CPT 2025 Professional Edition*. AMA; 2024. All rights reserved. CPT® is a registered trademark of the American Medical Association.
7. Centers for Medicare & Medicaid Services. Medicare claims processing manual. Chapter 12 - physicians/nonphysician practitioners. Accessed February 21, 2025. <https://www.cms.gov/regulations-and-guidance/guidance/manuals/downloads/clm104c12.pdf>

Please see Important Safety Information on pages **1** and **11-12** and accompanying full **Prescribing Information** for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

## SELECT IMPORTANT SAFETY INFORMATION (cont'd)

### CONTRAINDICATIONS

- SOLIRIS is contraindicated for initiation in patients with unresolved serious *Neisseria meningitidis* infection.

### WARNINGS AND PRECAUTIONS

#### Serious Meningococcal Infections

SOLIRIS, a complement inhibitor, increases a patient's susceptibility to serious, life-threatening, or fatal infections caused by meningococcal bacteria (septicemia and/or meningitis) in any serogroup, including non-groupable strains. Life-threatening and fatal meningococcal infections have occurred in both vaccinated and unvaccinated patients treated with complement inhibitors.

Revaccinate patients in accordance with ACIP recommendations considering the duration of therapy with SOLIRIS. Note that ACIP recommends an administration schedule in patients receiving complement inhibitors that differs from the administration schedule in the vaccine prescribing information.

If urgent SOLIRIS therapy is indicated in a patient who is not up to date with meningococcal vaccines according to ACIP recommendations, provide the patient with antibacterial drug prophylaxis and administer meningococcal vaccines as soon as possible. Various durations and regimens of antibacterial drug prophylaxis have been considered, but the optimal durations and drug regimens for prophylaxis and their efficacy have not been studied in unvaccinated or vaccinated patients receiving complement inhibitors, including SOLIRIS. The benefits and risks of treatment with SOLIRIS, as well as those associated with antibacterial drug prophylaxis in unvaccinated or vaccinated patients, must be considered against the known risks for serious infections caused by *Neisseria meningitidis*.

Vaccination does not eliminate the risk of serious meningococcal infections, despite development of antibodies following vaccination.

Closely monitor patients for early signs and symptoms of meningococcal infection and evaluate patients immediately if infection is suspected. Inform patients of these signs and symptoms and instruct patients to seek immediate medical care if these

signs and symptoms occur. Promptly treat known infections. Meningococcal infection may become rapidly life-threatening or fatal if not recognized and treated early. Consider interruption of SOLIRIS in patients who are undergoing treatment for serious meningococcal infection, depending on the risks of interrupting treatment in the disease being treated.

#### ULTOMIRIS and SOLIRIS REMS

Due to the risk of serious meningococcal infections, SOLIRIS is available only through a restricted program called ULTOMIRIS and SOLIRIS REMS.

Prescribers must enroll in the REMS, counsel patients about the risk of serious meningococcal infection, provide patients with REMS educational materials, assess patient vaccination status for meningococcal vaccines (against serogroups A, C, W, Y, and B) and vaccinate if needed according to current ACIP recommendations two weeks prior to the first dose of SOLIRIS. Antibacterial drug prophylaxis must be prescribed if treatment must be started urgently and the patient is not up to date with both meningococcal vaccines according to current ACIP recommendations at least two weeks prior to the first dose of SOLIRIS. Patients must receive counseling about the need to receive meningococcal vaccines and to take antibiotics as directed, the signs and symptoms of meningococcal infection, and be instructed to carry the Patient Safety Card with them at all times during and for 3 months following SOLIRIS treatment.

Further information is available at [www.UltSolREMS.com](http://www.UltSolREMS.com) or 1-888-765-4747.

#### Other Infections

Serious infections with *Neisseria* species (other than *Neisseria meningitidis*), including disseminated gonococcal infections, have been reported.

SOLIRIS blocks terminal complement activation; therefore, patients may have increased susceptibility to infections, especially with encapsulated bacteria, such as infections with *Neisseria meningitidis* but also *Streptococcus pneumoniae*, *Haemophilus influenzae*, and to a lesser extent, *Neisseria gonorrhoeae*.

Additionally, *Aspergillus* infections have occurred in immunocompromised and neutropenic patients. Patients receiving SOLIRIS are at increased risk for infections due to these organisms, even if they develop antibodies following vaccination.

Please see Important Safety Information on pages 1 and 11-12 and accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

## SELECT IMPORTANT SAFETY INFORMATION (cont'd)

### WARNINGS AND PRECAUTIONS (cont'd)

#### Thrombosis Prevention and Management

The effect of withdrawal of anticoagulant therapy during SOLIRIS treatment has not been established. Therefore, treatment with SOLIRIS should not alter anticoagulant management.

#### Infusion-Related Reactions

Administration of SOLIRIS may result in infusion-related reactions, including anaphylaxis or other hypersensitivity reactions. In clinical trials, no patients experienced an infusion-related reaction which required discontinuation of SOLIRIS. Interrupt SOLIRIS infusion and institute appropriate supportive measures if signs of cardiovascular instability or respiratory compromise occur.

### ADVERSE REACTIONS

The most frequently reported adverse reactions in the NMOSD placebo-controlled trial ( $\geq 10\%$ ) were: upper respiratory infection, nasopharyngitis, diarrhea, back pain, dizziness, influenza, arthralgia, pharyngitis, and contusion.

### DRUG INTERACTIONS

#### Plasmapheresis, Plasma Exchange, or Fresh Frozen Plasma Infusion

Concomitant use of SOLIRIS with plasma exchange (PE), plasmapheresis (PP), or fresh frozen plasma infusion (PE/PI) treatment can reduce serum eculizumab concentrations and requires a supplemental dose of SOLIRIS.

#### Neonatal Fc Receptor Blockers

Concomitant use of SOLIRIS with neonatal Fc receptor (FcRn) blockers may lower systemic exposures and reduce effectiveness of SOLIRIS. Closely monitor for reduced effectiveness of SOLIRIS.

To report SUSPECTED ADVERSE REACTIONS contact Alexion Pharmaceuticals, Inc. at 1-844-259-6783 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

Please see accompanying full [Prescribing Information](#) for SOLIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.